

In the Claims

1 (currently amended). ~~A~~ An isolated polypeptide, which polypeptide that:

- (i) comprises or consists of the amino acid sequence as recited in SEQ ID NO:2;
- (ii) is a fragment of SEQ ID NO:2 thereof that is ~~an interferon-gamma-like secreted protein of the four helical bundle cytokine fold, or having an antigenic determinant in common with polypeptides of (i);~~ or
- (iii) ~~is a functional equivalent of (i) or (ii)~~ has greater than 90% sequence identity with the polypeptide of SEQ ID NO: 2 or a fragment thereof;

wherein said isolated polypeptide or fragment has anti-viral activity, antiproliferative activity on cancer cells, the ability to increase levels of TNF- α , IL-2, IL-4, IFN- γ , aspartate aminotransferase (ASAT) or alanine aminotransferase (ALAT) *in vivo*, or the ability to induce IFN- γ secretion by concanavalin A or phytohemagglutinin stimulated peripheral blood mononuclear cells.

2-5 (canceled).

6 (currently amended). ~~The fragment or functional equivalent of claim 4~~ isolated polypeptide according claim 1, wherein said polypeptide has greater than 95% sequence identity with the polypeptide of SEQ ID NO: 2 or a fragment thereof and has anti-viral activity, antiproliferative activity on cancer cells, the ability to increase levels of TNF- α , IL-2, IL-4, IFN- γ , aspartate aminotransferase (ASAT) or alanine aminotransferase (ALAT) *in vivo*, or the ability to induce IFN- γ secretion by concanavalin A or phytohemagglutinin stimulated peripheral blood mononuclear cells
there is greater than 95% sequence identity.

7 (currently amended). ~~The fragment or functional equivalent of claim 4~~ isolated polypeptide according claim 1, wherein said polypeptide has greater than 98% sequence identity with the polypeptide of SEQ ID NO: 2 or a fragment thereof and has anti-viral activity, antiproliferative activity on cancer cells, the ability to increase levels of TNF- α , IL-2, IL-4, IFN- γ , aspartate

aminotransferase (ASAT) or alanine aminotransferase (ALAT) *in vivo*, or the ability to induce IFN- γ secretion by concanavalin A or phytohemagglutinin stimulated peripheral blood mononuclear cells; wherein there is greater than 98% sequence identity.

8 (currently amended). ~~The fragment or functional equivalent of claim 4 isolated polypeptide according to claim 1, wherein said polypeptide has greater than 99% sequence identity with the polypeptide of SEQ ID NO: 2 or a fragment thereof and has anti-viral activity, antiproliferative activity on cancer cells, the ability to increase levels of TNF- α , IL-2, IL-4, IFN- γ , aspartate aminotransferase (ASAT) or alanine aminotransferase (ALAT) *in vivo*, or the ability to induce IFN- γ secretion by concanavalin A or phytohemagglutinin stimulated peripheral blood mononuclear cells~~ wherein there is greater than 99% sequence identity.

9-50 (canceled).

51 (new). The isolated polypeptide according to claim 1, wherein said polypeptide comprises the amino acid sequence as recited in SEQ ID NO: 2.

52 (new). The isolated polypeptide according to claim 1, wherein said polypeptide consists of the amino acid sequence as recited in SEQ ID NO: 2.

53 (new). The isolated polypeptide according to claim 1, wherein said polypeptide is a fragment of SEQ ID NO: 2 and has anti-viral activity, antiproliferative activity on cancer cells, the ability to increase levels of TNF- α , IL-2, IL-4, IFN- γ , aspartate aminotransferase (ASAT) or alanine aminotransferase (ALAT) *in vivo*, or the ability to induce IFN- γ secretion by concanavalin A or phytohemagglutinin stimulated peripheral blood mononuclear cells.

54 (new). The isolated polypeptide according to claim 1, wherein said polypeptide is a fragment of a polypeptide having greater than 90% sequence identity to SEQ ID NO: 2 and said fragment has anti-viral activity, antiproliferative activity on cancer cells, the ability to increase levels

of TNF- α , IL-2, IL-4, IFN- γ , aspartate aminotransferase (ASAT) or alanine aminotransferase (ALAT) *in vivo*, or the ability to induce IFN- γ secretion by concanavalin A or phytohemagglutinin stimulated peripheral blood mononuclear cells.

- 55 (new). A composition comprising a pharmaceutical carrier and a polypeptide that:
- (i) comprises or consists of the amino acid sequence as recited in SEQ ID NO:2;
 - (ii) is a fragment of SEQ ID NO:2; or
 - (iii) has greater than 90% sequence identity with the polypeptide of SEQ ID NO: 2 or a fragment thereof;

wherein said isolated polypeptide or fragment has anti-viral activity, antiproliferative activity on cancer cells, the ability to increase levels of TNF- α , IL-2, IL-4, IFN- γ , aspartate aminotransferase (ASAT) or alanine aminotransferase (ALAT) *in vivo*, or the ability to induce IFN- γ secretion by concanavalin A or phytohemagglutinin stimulated peripheral blood mononuclear cells.

56 (new). The composition according to claim 55, wherein said polypeptide has greater than 95% sequence identity with the polypeptide of SEQ ID NO: 2 or a fragment thereof and has anti-viral activity, antiproliferative activity on cancer cells, the ability to increase levels of TNF- α , IL-2, IL-4, IFN- γ , aspartate aminotransferase (ASAT) or alanine aminotransferase (ALAT) *in vivo*, or the ability to induce IFN- γ secretion by concanavalin A or phytohemagglutinin stimulated peripheral blood mononuclear cells.

57 (new). The composition according to claim 55, wherein said polypeptide has greater than 98% sequence identity with the polypeptide of SEQ ID NO: 2 or a fragment thereof and has anti-viral activity, antiproliferative activity on cancer cells, the ability to increase levels of TNF- α , IL-2, IL-4, IFN- γ , aspartate aminotransferase (ASAT) or alanine aminotransferase (ALAT) *in vivo*, or the ability to induce IFN- γ secretion by concanavalin A or phytohemagglutinin stimulated peripheral blood mononuclear cells.

58 (new). The composition according to claim 55, wherein said polypeptide has greater than 99% sequence identity with the polypeptide of SEQ ID NO: 2 or a fragment thereof and has anti-viral activity, antiproliferative activity on cancer cells, the ability to increase levels of TNF- α , IL-2, IL-4, IFN- γ , aspartate aminotransferase (ASAT) or alanine aminotransferase (ALAT) *in vivo*, or the ability to induce IFN- γ secretion by concanavalin A or phytohemagglutinin stimulated peripheral blood mononuclear cells.

59 (new). The composition according to claim 55, wherein said polypeptide comprises the amino acid sequence as recited in SEQ ID NO: 2.

60 (new). The composition according to claim 55, wherein said polypeptide consists of the amino acid sequence as recited in SEQ ID NO: 2.

61 (new). The composition according to claim 55, wherein said polypeptide is a fragment of SEQ ID NO: 2 and has anti-viral activity, antiproliferative activity on cancer cells, the ability to increase levels of TNF- α , IL-2, IL-4, IFN- γ , aspartate aminotransferase (ASAT) or alanine aminotransferase (ALAT) *in vivo*, or the ability to induce IFN- γ secretion by concanavalin A or phytohemagglutinin stimulated peripheral blood mononuclear cells.

62 (new). The composition according to claim 55, wherein said polypeptide is a fragment of a polypeptide having greater than 90% sequence identity to SEQ ID NO: 2 and said fragment has anti-viral activity, antiproliferative activity on cancer cells, the ability to increase levels of TNF- α , IL-2, IL-4, IFN- γ , aspartate aminotransferase (ASAT) or alanine aminotransferase (ALAT) *in vivo*, or the ability to induce IFN- γ secretion by concanavalin A or phytohemagglutinin stimulated peripheral blood mononuclear cells.